


FOREIGN TECHNOLOGY DIVISION



CALCULATION OF THE THERMODYNAMIC PROPERTIES
OF DISSOCIATING NITROGEN TETROXIDE
TAKING NONIDEALITY INTO ACCOUNT

by

V. B. Nesterenko, M. A. Bazhin, and V. P. Bubnov



Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

AD 702 989

EDITED TRANSLATION

CALCULATION OF THE THERMODYNAMIC PROPERTIES OF
DISSOCIATING NITROGEN TETROXIDE TAKING NONIDEALITY
INTO ACCOUNT

By: V. B. Nesterenko, M. A. Bazhin, and
V. P. Bubnov

English pages: 7

Source: AN BSSR. Vestsi. Seryya Fizika-Tekhnichnykh
Navuk (Academy of Sciences of the Belorussian
SSR. News. Series in the Physical and
Technical Sciences), No. 3, 1966, pp. 20-24.

Translated by: L. Heenan/TDBRS-3

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.


PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

CALCULATION OF THE THERMODYNAMIC PROPERTIES
OF DISSOCIATING NITROGEN TETROXIDE
TAKING NONIDEALITY INTO ACCOUNT

V. B. Nesterenko, M. A. Bazhin, and V. P. Bubnov

Recently there has been considerable interest in chemically reactive systems as the most effective working substances [1, 2] for power plants. Nitrogen tetroxide, whose dissociation occurs in the temperature range 290-1500°K,



is the most frequently studied of a large class of chemically reactive systems.

In order to perform a thermodynamic analysis of the cycles of power plants on a dissociating working substance and to explain the advantages of such systems, it is necessary to know the enthalpy and entropy in a wide range of temperatures and pressures.

At the present time there is virtually no experimental data on the enthalpy or the heat capacity of the dissociating system $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2 \rightleftharpoons 2\text{NO} + \text{O}_2$. It is also impossible to plot thermodynamic diagrams on the basis of experimental P-V-T data since only one work (by W. G. Schlinger and B. H. Sage [3]) has been published on the determination of P-V-T data for N_2O_4 in the gaseous phase with a temperature range of 294-440°K and pressures of 1-140 atm (abs).

Based on this, we have attempted to calculate the thermodynamic functions (enthalpy and entropy) using the common thermodynamic relationships and taking into account nonideality for chemically reactive nitrogen tetroxide.

Determining composition. In the thermal dissociation of nitrogen tetroxide



a mixture is formed whose composition changes with a change in pressure and temperature. In order to determine the equilibrium composition of the reactive mixture, the equilibrium constant K_p is used, which, in the case of reaction (1), can be represented in the form

$$K_{p_1} = \frac{4a_1^2(1-a_2)^2}{(1-a_1)(1+a_1+a_1a_2)} P, \quad (2)$$

$$K_{p_2} = \frac{a_1a_2^2}{(1-a_2)^2(1+a_1+a_1a_2)} P. \quad (3)$$

From equations (2) and (3) it follows that in order to determine the molar content of the components in the mixture, it is necessary to know for each value of P and T the size of the equilibrium constants K_{p_1} and K_{p_2} .

When we can apply the principle of volume additivity (Amagat's law) to the thermodynamic system, quantities K_{p_j} can be calculated as

$$K_{p_j} = \frac{K_{p_0j}}{K_{v_j}}, \quad (4)$$

where quantities K_{v_j} are found from the volatility data of the mixture's components. For the examined reaction (1) quantities K_{v_j} are determined by the relationship

$$K_{v_1} = \frac{v_{\text{N}_2\text{O}_4}}{v_{\text{NO}_2}^2}; K_{v_2} = \frac{v_{\text{NO}_2}^2}{v_{\text{NO}}^2 v_{\text{O}_2}}, \quad (5)$$

in which the coefficient of activity $\gamma_j = \frac{f_j}{P}$.

In view of the fact that the components N_2O_4 and NO_2 do not exist in pure form, their coefficient of activity was determined on the basis of the experimental P-V-T data of W. G. Schlinger and B. H. Sage [3] according to formula

$$\ln \gamma = \int_0^P \left(\frac{V}{RT} - \frac{1}{P} \right) dP. \quad (6)$$

The values of the activity coefficients for components NO and O_2 were obtained from the relationship $\gamma = f(\tau, \pi)$ [4].

Based on the obtained data for the activity coefficient, the values of K_j were calculated in the temperature range 300-1500°K and pressure range 1-150 atm (abs).

Since the polynomial of the equilibrium constant obtained by Bodenstein [5] is limited by a temperature range of ~650°K in the first stage of the reaction, we used values for K_{p01} based on a calculation of the isobaric-isothermal potentials for the N_2O_4 and NO_2 components according to spectroscopic and molecular data [6, 7] and the more precisely defined thermal effect of the hypothetical reaction of a nitrogen tetroxide association, according to the experimental data of Bodenstein and Bouise [8], Verhock and Daniels [9], and Wourtsel [10].

The values for the polynomial of the equilibrium constant K_{p20} were taken from the work of Bodenstein and Lindner [8].

On the basis of the values obtained for K_{p10} and K_{p20} we calculated the equilibrium composition of the chemically reactive system



The thermodynamic functions (enthalpy and entropy) necessary for plotting I-S and T-S diagrams were calculated on the basis of thermodynamic relationships, the thermodynamic theory of empirical

corrections, and generalized tables.

As Hougen, Watson, and Ragatz have shown [11], the molar entropy for real gas can be expressed as

$$S = S_{\text{ideal}} p_0^{-R} \ln (P/P_0) - (S_{\text{ideal}} - S)_{p,T}, \quad (8)$$

where the correction for nonideality in terms of relative parameters is calculated thus:

$$(S_{\text{га}} - S)_{p,T} = -R \int_0^1 \frac{(1-z)}{\pi} d\pi + R \tau \int_0^1 \left(\frac{\partial z}{\partial \tau} \right) \frac{d\pi}{\pi} \quad (9)$$

[ид = ideal]

Hence the value of entropy for the mixture, relative to an initial mole of the nondissociated substance N_2O_4 , has the form:

$$\begin{aligned} S_{\text{см}} = & \frac{1}{M_{N_2O_4}} \left\{ \sum_{i=1}^m \left(\frac{n_i}{n_{N_2O_4}} S_{\text{ид}i} \right) - \right. \\ & - R \sum_{i=1}^m \left[\frac{n_i}{n_{N_2O_4}} \ln \left(\frac{n_i}{N} \right) - \right. \\ & \left. \left. - \frac{N}{n_{N_2O_4}} R \ln \left(\frac{P}{P_0} \right) - \right. \right. \\ & \left. \left. - \sum_{i=1}^m \left[\frac{n_i}{n_{N_2O_4}} (S_{\text{га}i} - S)_{p,T} \right] \right\} \right. \\ & \text{[см = mixture]} \\ & \text{[ид = ideal]} \end{aligned} \quad (10)$$

In equation (10) the first term represents the entropy of each ideal component in a gaseous phase (8); the next term is the entropy of the mixture; the following term takes into account the variation in entropy from standard pressure to desirable pressure; the last term is the correction for the deviation from ideal gas behavior.

For the examined system the entropy of the mixture is written by analogy with equation (10)

$$S_{\text{cm}} = \frac{1}{M_{\text{N}_2\text{O}_4}} \left\{ \left[(1-a_1) S_{\text{N}_2\text{O}_4} + 2a_1(1-a_2) S_{\text{NO}_2} + 2a_1a_2 \left(S_{\text{NO}} + \frac{1}{2} S_{\text{O}_2} \right) \right] - R \left[(1-a_1) \ln(1-a_1) + 2a_1a_2 \ln 2a_1a_2 + a_1a_2 \ln a_1a_2 + \right. \right. \\ \left. \left. + 2a_1(1-a_2) \ln 2a_1(1-a_2) - (1+a_1+a_1a_2) \ln(1+a_1+a_1a_2) \right] - \right. \\ \left. - [(1-a_1)(S_{\text{N}_2} - S)_{\text{N}_2\text{O}_4} + 2a_1(1-a_2)(S_{\text{N}_2} - S)_{\text{NO}_2} + \right. \\ \left. + 2a_1a_2(S_{\text{N}_2} - S)_{\text{NO}} + a_1a_2(S_{\text{N}_2} - S)_{\text{O}_2}] \right\}. \quad (11)$$

The enthalpy of the mixture is written as

$$I_{\text{cm}} = \sum_{i=1}^n \frac{n_i I_i}{M_{\text{N}_2\text{O}_4}}. \quad (12)$$

In view of the fact that we previously presented a method for calculating the volatility factor, it is advisable to express the correction for nonideality in equations (9) and (12) in terms of the activity coefficient:

$$S_i = S_{0i} - \left[RT \left(\frac{\partial \ln \gamma_i}{\partial T} \right)_P + R \ln \gamma_i \frac{P}{P_0} \right], \quad (13)$$

$$I_i = I_{0i} - RT^2 \left(\frac{\partial \ln \gamma_i}{\partial T} \right)_P, \quad (14)$$

$$\left. \begin{aligned} S_{0i} &= s_{0i} + S_i^0 \\ I_{0i} &= i_{0i} + I_i^0 \end{aligned} \right\}. \quad (15)$$

The value of quantities i_{0j} , s_{0j} , S_j^0 , I_j^0 is taken from reference [12].

On the basis of the obtained relationships the values of enthalpy and entropy in the temperature range 300-1500°K and pressure range 1-150 atm (abs) were computed on the Minsk-2 computer at the Computer Center of the BSSR Academy of Sciences.

The values of enthalpy and entropy on the saturation line were obtained from the experimental P-V-T data of Reamer and Sage [13] on the saturation line. The I-S and T-S diagrams which the authors obtained for the dissociating system $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2 \rightleftharpoons 2\text{NO} + \text{O}_2$ are presented in Figs. 1 and 2.




Fig. 1. I-S diagram of the dissociating system $N_2O_4 \rightleftharpoons 2NO_2 \rightleftharpoons 2NO + O_2$.




Fig. 2. T-S diagram of the dissociating system $N_2O_4 \rightleftharpoons 2NO_2 \rightleftharpoons 2NO + O_2$.

Designations

P - total pressure of mixture in kg/cm^2 ; α_1, α_2 - degree of dissociation of first and second stages in the reaction; $K_{p_{0j}}$ - equilibrium constant as a function of temperature only; f_j - volatility of the j -th component taken at pressure P of mixture; n_j - content of

moles of j-th component in mixture; M - molecular weight; π - relative pressure; τ - relative temperature; T - temperature in $^{\circ}\text{K}$; z - compressibility factor; P_0 - initial pressure; i_{0j} , s_{0j} - relative enthalpy and entropy; i_j^0 , s_j^0 - enthalpy and entropy under standard conditions, $N = 1 + \alpha_1 + \alpha_1\alpha_2$.

Literature

1. Laytkhill, M. Voprosy raketnoy tekhniki (Problems in rocketry), No. 5, 1957.
2. Stai, D. F., F. Bizjak, and S. E. Stephanou. J. Spacecraft, 2, No. 5, 742-745, 1965.
3. Schlinger, W. G., and B. H. Sage. Ind. and Engineering Chem., 42, No. 10, 1952.
4. Karapet'yants, M. Kh. Khimicheskaya termodinamika (Chemical thermodynamics), 1953.
5. Bodenstein, M., and Boüse. Phys. Chem., 100, 1922.
6. Smith, D. K., and K. Hedberg. J. Ch. Ph., 25, No. 6, 1956.
7. Hisatsune, J. C. Advances in molecular spectrosc., 3, 1962.
8. Bodenstein, M., and Lindner. Zeit. Chem., 100, 1922.
9. Verhock, F. H., and F. J. Daniels. Amer. Ch. Soc., 53, 1950, 1931.
10. Wourtsel, M. Comptes Rendus (Reports) 169, 1937, 1919.
11. Hougen, O. A., K. M. Watson, and R. A. Ragatz. Chemical Process Principles, 2, 1959.
12. Nesterenko, V. B., and V. P. Bubnov. Vestsi AN BSSR, ser. fiz.-tekhn., No. 1, 1966.
13. Reamer, H. H., and B. H. Sage. Ind. Eng. Chem., 44, 1952.

I.Ya.E., BSSR Academy of Sciences

Received
23 March 1966

DATA HANDLING PAGE				
01-ACCESSION NO. 98-DOCUMENT LOC		39-TOPIC TAGS		
TP9001870		nitrogen tetroxide, nitrogen tetroxide dissociation, entropy, enthalpy, thermodynamic function		
09-TITLE CALCULATION OF THE THERMODYNAMIC PROPERTIES OF DISSOCIATING NITROGEN TETROXIDE TAKING NONIDEALITY INTO ACCOUNT -U-				
47-SUBJECT AREA				
20				
42-AUTHOR CL JTHORS NESTERENKO, V. B.; 16-BAZHIN, M. A.; 16-BUBNOV, V. P.				10-DATE OF INFO -----66
43-SOURCE AN BSSR. VESTSI. SERYYA FIZIKA TEKHNICHNYKH NAVUK (RUSSIAN)				68-DOCUMENT NO. HT-23-303-69
				69-PROJECT NO. 60401
63-SECURITY AND DOWNGRADING INFORMATION			64-CONTROL MARKINGS	97-HEADER CLASN
UNCL. 0			NONE	UNCL
76-REEL/FRAME NO.	77-SUPERSEDES	78-CHANGES	40-GEOGRAPHICAL AREA	NO. OF PAGES
1890 1548			UR	7
CONTRACT NO. F33057-68-D-0865-P002	X REF ACC. NO. 65-AP6033066	PUBLISHING DATE 94-	TYPE PRODUCT TRANSLATION	REVISION FREQ NONE
STEP NO. 02-UR/0201/66/000/003/0020/0024				
ABSTRACT				
<p>(U) This study was undertaken because of the lack of experimental data on the enthalpy of dissociating nitrogen tetroxide. At calculation of the entropy (S) and enthalpy (I) of dissociating N_2O_4 was performed in the temperature range 300-1500K and 1-140 atm range, taking into account deviation of the reaching N_2O_4 from ideal behavior. The calculation was carried out on the basis of general thermodynamic functions of the thermodynamic theory of empirical corrections and of generalized tables. The calculated S and I values were used for plotting I-S and T-S diagrams. Orig. art. has: 2 figures.</p>				